

---

# Neurology

## Seven Things Clinicians and Patients Should Question

by  
Canadian Neurological Society  
Last updated: July 2022



### 1 **Don't routinely perform brain imaging after acute seizure in patients with established epilepsy.**

Typically, epilepsy patients have brain imaging at the time of diagnosis to investigate for a structural cause for seizures. Seizures in these patients are unlikely a result of new structural changes. Neuroimaging can be considered in patients with longstanding epilepsy without prior imaging studies, or in patients who are candidates for neurosurgery with seizures refractory to medical management.

### 2 **Don't treat women of childbearing potential with Valproic acid if other effective treatments are available.**

Valproic acid is teratogenic for a developing fetus and should be avoided. Valproic acid increases the risk of diminished IQ as well as major congenital malformations including neural tube defects and cardiovascular abnormalities for a developing fetus. If valproic acid treatment is deemed necessary, patients should be counselled regarding contraception and the risks of conception. In this case, the lowest effective dose should be used.

### 3 **Don't choose opioids or cannabinoids as the first choice of treatment for neuropathic pain.**

Opioids and cannabinoids have weak or inconclusive evidence in effective treatment of neuropathic pain. The well documented risks of opioid and cannabinoids include nausea, sleepiness, impairment, dependence, and development of substance use disorders. With impairment comes further risks to oneself and others in altered judgement in the workplace or while operating a vehicle. Opioids come with an additional risk of decreased respiratory drive and fatality with overdose. Neuropathic pain can be treated effectively using agents with demonstrated efficacy and significantly less risks compared to opioids and cannabinoids.

### 4 **Don't order neuroimaging or EEG in asymptomatic patients in the emergency department with syncope and a normal neurological evaluation.**

Syncope, and pre-syncope with observed convulsions are very common and frequently investigated in the emergency department. Typical syncope with a normal examination requires minimal investigation. Neuroimaging and EEG will not help in determining the etiology or management of patients with typical syncope in the absence of focal neurologic symptoms or findings on examination.

### 5 **Don't use opioids for treatment of migraines.**

Opioids are not adequate for pain control for patients with migraines. The risk for harm, including impairment, dependence, tolerance, medication overuse headaches, and opioid use disorder with opioids is greater than the documented benefit. Additionally, opioids may worsen nausea and vomiting associated with the migraine. Prescription opioids for migraines would have minimal to no benefit with the excess of risk, and contribute to the opioid crisis.

---

## How the list was created

The Canadian Neurological Society (CNS) formed a Choosing Wisely task force in August 2018. The American Board of Internal Medicine Choosing Wisely Neuroscience recommendations were reviewed for their relevance to Canadian neurology as well as recommendations from the American Epilepsy Society, American Association of Neuromuscular and Electrodiagnostic Medicine, American Academy of Nursing, American College of Emergency Physicians, American Geriatrics Society, American Academy of Neurology, and American Academy of Pediatrics. Ten potential recommendations were initially selected and were sent to CNS members in a ranking survey from which a top five list was chosen. These recommendations were presented at the Canadian Neurosciences Federation National Congress in June 2019 which were well received. The list was then sent to Choosing Wisely Canada for review and finalization.

---

## Sources

- 1** Cendes F, et al. Neuroimaging of epilepsy. Handbook of Clinical Neurology. 2016;136:985-1014. [PMID: 27430454](#).  
Commission on Neuroimaging of the International League Against Epilepsy. Recommendations for neuroimaging of patients with epilepsy. Epilepsia. 1997 Nov;38(11):1255-6. [PMID: 9579930](#).  
Epilepsy Implementation Task Force. [Provincial Guidelines for the Management of Epilepsy in Adults and Children](#). Critical Care Services Ontario. January 2015. [Internet].  
Harden CL, et al. Reassessment: Neuroimaging in the emergency patient presenting with seizure (an evidence-based review). Report of the Therapeutics and Technology Assessment Subcommittee of the American Academy of Neurology. Neurology. 2007 Oct 30;69(18):1772-80. [PMID: 17967993](#).  
Lapalme-Remis S, and Cascino GD. Imaging for Adults with Seizures and Epilepsy. Continuum (Minneapolis, Minn). 2016 Oct;22(5, Neuroimaging):1451-1479. [PMID: 27740984](#).
- 2** Bansal R, et al. [Maternal and neonatal complications during pregnancy in women with epilepsy](#). International Journal of Epilepsy. 2016 July;3(2):80-85.  
Campbell E, et al. Malformation risks of antiepileptic drug monotherapies in pregnancy: updated results from the UK and Ireland Epilepsy and Pregnancy Registers. J Neurol Neurosurg Psychiatry. 2014 Sep;85(9):1029-34. [PMID: 24444855](#).  
Hernández-Díaz S, et al for the North American AED Pregnancy Registry. Comparative safety of antiepileptic drugs during pregnancy. Neurology. 2012 May;78(21):1692-9. [PMID: 22551726](#).  
Voinescu PE and Pennell PB. Management of epilepsy during pregnancy. Expert Rev Neurother. 2015 Oct;15(10):1171-87. [PMID: 26416395](#).
- 3** Moulin DE, et al. Pharmacological management of chronic neuropathic pain: Revised consensus statement from the Canadian Pain Society. Pain Res Manag. 2014 Nov-Dec;19(6):328-35. [PMID: 25479151](#).  
Finnerup NB, et al. Pharmacotherapy for neuropathic pain in adults: a systematic review and meta-analysis. Lancet Neurol. 2015 Feb;14(2):162-73. [PMID: 25575710](#).
- 4** Idil H and Yilmaz K. Diagnostic yield of neuroimaging in syncope patients without high-risk symptoms indicating neurological syncope. Am J Emerg Med. 2018 May 16. pii: S0735-6757(18)30410-8. [PMID: 29802003](#).  
Sanatani S et al. Canadian Cardiovascular Society and Canadian Pediatric Cardiology Association Position Statement on the Approach to Syncope in the Pediatric Patient. Can J Cardiol. Feb 2017;33(2):189-198. [PMID: 27838109](#).  
Shenk MER and Anilkumar A. [Syncope in Children: Etiology, Positive and Negative Predictors, and Utilization and Utility of Diagnostic Testing](#). Neurology. Neurology. Apr 2017;88(16):P3.204.
- 5** Tepper S & Spears RC. Acute treatment of migraine. Neurologic Clinics. 2009 May;27(2):417-27. [PMID: 19289223](#).  
Worthington I, et al. Canadian Headache Society Guideline: Acute Drug Therapy for Migraine Headache. Canadian Journal of Neurological Sciences / Journal Canadien Des Sciences Neurologiques. Sept 2013;40(S3):S1-S3. [PMID: 23968886](#).

---

## About the Canadian Neurological Society

The Canadian Neurological Society was established in 1948 and represented both neurologists and neurosurgeons. In 1965, the original CNS was dissolved and two new societies were created to represent the two distinct groups.

Today, the Canadian Neurological Society has approximately 540 members and represents neurologists and neurology residents in Canada. The CNS Board of Directors manages the business, initiatives and finances of the CNS. The board also appoints CNS member representatives to sit on various CNSF committees, including the Professional Development Committee and the Scientific Program Committee. The CNS President and Vice-President also serve on the Board of Directors of the Canadian Neurological Sciences Federation (CNSF).



---

## About Choosing Wisely Canada

Choosing Wisely Canada is the national voice for reducing unnecessary tests and treatments in health care. One of its important functions is to help clinicians and patients engage in conversations that lead to smart and effective care choices.

🌐 [ChoosingWiselyCanada.org](http://ChoosingWiselyCanada.org) | ✉ [info@ChoosingWiselyCanada.org](mailto:info@ChoosingWiselyCanada.org) | 🐦 [@ChooseWiselyCA](https://twitter.com/ChooseWiselyCA) | 📘 [/ChoosingWiselyCanada](https://www.facebook.com/ChoosingWiselyCanada)